arXiv:1702.02614v1 [math.GT] 8 Feb 2017

THE NUMBER OF NON-NEGATIVE CURVATURE TRIANGULATIONS OF S^2

PHILIP ENGEL[†] AND PETER SMILLIE

ABSTRACT. A triangulation of S^2 has *non-negative curvature* if every vertex has at most six triangles adjacent to it. Thurston showed that the non-negative curvature triangulations of S^2 correspond to orbits of vectors of positive norm in an Eisenstein lattice $\Lambda \subset \mathbb{C}^{1,9}$. By integrating the Siegel theta function associated to Λ, we show that the appropriately weighted number of triangulations of S^2 with $2n$ triangles is

$$\frac{809}{2612138803200}\sigma_9(n),$$

where $\sigma_k(n) = \sum_{d|n} d^k$ is the kth divisor function.

1. INTRODUCTION

Triangulations are frequently used to model a random surface. An essential ingredient in such a model is the number of triangulations with a given number of triangles. For instance, to model a random metric on a surface, one can declare each triangle to be isometric to a fixed equilateral triangle. On the sphere, it is natural to restrict to triangulations whose associated metric is non-negatively curved. We say that a triangulation of the sphere has *non-negative curvature* if every vertex is adjacent to six or fewer triangles. Such triangulations arise as the dual complex of a degeneration of $K3$ surfaces in (-1)-form [10]. Similarly, the dual complex of a spherical fullerene, or buckyball, is a non-negative curvature triangulation of S^2.

Non-negative curvature is a very strong restriction which changes the nature of the counting problem. Whereas the total number of triangulations grows exponentially in the number of triangles, Thurston [17] showed that the number of non-negative curvature triangulations with up to $2n$ triangles is $O(n^{10})$. More specifically, by interpreting them as integral points in a moduli space of metric surfaces, he proved that non-negative curvature triangulations with $2n$ triangles correspond to $U(\Lambda)$-orbits of vectors of square norm $2n$ in a Hermitian lattice $\Lambda \subset \mathbb{C}^{1,9}$.

In Section 2, we begin by summarizing Thurston's work. In Theorem 2.3, we prove a formula for the weighted number of non-negative curvature triangulations. The main input for this result is Corollary 3.5 of Theorem 3.4, which states that the weighted count of $U(\Lambda)$-orbits of vectors of square norm $2n$ is the nth Fourier coefficient of a scalar multiple of the Eisenstein

[†]Research partially supported by NSF grant DMS-1502585.

series $E_{10}(\tau)$. In Appendix A, we explicitly compute the Hermitian pairing on the lattice Λ. In Appendix B, we sketch how our method gives a formula for the number of non-negative curvature square-tilings, or squarulations, of the sphere.

Acknowledgements: We would like to thank Simion Filip, Karsten Gimre, and Curtis McMullen for their suggestions and conversation.

2. Counting triangulations of S^2

A *flat cone sphere* is a sphere with a flat metric away from a finite set of points $\{p_1, \dots, p_n\}$, which near each point is isometric to a convex Euclidean cone. In 1942, Alexandrov [1] proved that a flat cone sphere is isometric to a unique convex polyhedron in $\mathbb{R}^3$. Near each singularity p_i the angle of the cone is $2\pi - \alpha_i$ for some $\alpha_i \in (0, 2\pi)$. We call α_i the *cone angle deficit*. The Gauss-Bonnet theorem implies that

$$\sum \alpha_i = 4\pi.$$

Thurston [17] studied the moduli space $\mathcal{M}_{\alpha_1,\dots,\alpha_n}$ of all flat cone spheres modulo scaling with specified cone angle deficits at n unmarked points. Using local period maps to complex hyperbolic space

$$\mathbb{CH}^{n-3} := \mathbb{P}\{v \in \mathbb{C}^{1,n-3} \,\big|\, v \cdot v > 0\},$$

Thurston showed that $\mathcal{M}_{\alpha_1,\dots,\alpha_n}$ is a complex hyperbolic orbifold. Such moduli spaces were studied earlier by Deligne and Mostow [6], see also Looijenga's survey paper [11], in the context of hypergeometric functions and Lauricella differentials, i.e. differentials on $\mathbb{P}^1$ of the form

$$\eta = (z - z_1)^{-r_1} \dots (z - z_{n-1})^{-r_{n-1}}\, dz$$

where $r_i \in (0, 1)$. When $\sum r_i \in (1, 2)$, such a differential induces on $\mathbb{P}^1$ the structure of a flat cone sphere by using $\int_0^z \eta$ as a local coordinate.

The moduli space $\mathcal{M}_{\alpha_1,\dots,\alpha_n}$ is generally metrically incomplete because the cone singularities can collide, in which case the cone angle adds. Therefore Thurston considered the metric completion $\overline{\mathcal{M}}_{\alpha_1,\dots,\alpha_n}$, which is stratified by moduli spaces of flat cone spheres in which some collections of singularities have coalesced. One natural question is whether the completion is still a complex hyperbolic orbifold. If so, $\overline{\mathcal{M}}_{\alpha_1,\dots,\alpha_n}$ is isometric to the quotient of $\mathbb{CH}^{n-3}$ by the monodromy group $\Gamma \subset U(1, n)$ of $\mathcal{M}_{\alpha_1,\dots,\alpha_n}$, because complete metric orbifolds of nonpositive curvature are global quotients of manifolds.

The completion $\overline{\mathcal{M}}_{\alpha_1,\dots,\alpha_n}$ is always an orbifold when $n = 3$, since the moduli space is a point. For $n = 4$, there are infinitely many choices of α_i for which $\overline{\mathcal{M}}_{\alpha_1,\alpha_2,\alpha_3,\alpha_4}$ is a complex hyperbolic orbifold. In fact, every quotient by a triangle group $\Gamma \subset U(1, 1)$ acting on $\mathbb{CH}^1 = \mathbb{H}^2$ appears as a moduli space of tetrahedra. For $n = 5$, Picard [14] and his student LeVavasseur answered this question, in the guise of determining when the

monodromy group of the hypergeometric function

$$F(x,y) = \int_1^\infty z^{-r_1}(z-1)^{-r_2}(z-x)^{-r_3}(z-y)^{-r_4}\,dz$$

is discrete. For $n \geq 5$, Mostow [13] and later Thurston enumerated all 94 values of the α_i for which the moduli space is an orbifold.

The highest dimensional completed moduli space of flat cone spheres which is an orbifold corresponds to $n = 12$ and $\alpha_i = \pi/3$ for all i. See [15] for a more detailed treatment of this case. There is a stratification

$$\Gamma\backslash\mathbb{CH}^9 = \coprod_{\substack{\sum \mu_i = 12 \\ \mu_i \in \{1,2,3,4,5\}}} \mathcal{M}_{\mu_1\pi/3,\,\ldots,\,\mu_n\pi/3}.$$

Of the 78 completed moduli spaces of flat cone spheres with $n \geq 5$ which are orbifold quotients of $\mathbb{CH}^{n-3}$ by arithmetic groups, 34 are strata in the above example. The total number of strata is 45.

One special feature of this example is that Γ preserves a lattice $\Lambda \subset \mathbb{C}^{1,9}$ which has geometric significance in the moduli interpretation—the flat cone spheres which admit a triangulation into metrically equilateral triangles are the projectivization $\mathbb{P}\Lambda \cap \mathbb{CH}^9$ of this lattice. Conversely, a non-negative curvature triangulation of S^2 admits the structure of a flat cone sphere lying in one of the above strata by declaring each triangle to be metrically equilateral. In fact, Thurston, cf. Theorem 0.1 of [17], showed there is a correspondence

$$\{\text{Oriented non-negative curvature triangulations of } S^2\} \longleftrightarrow \Gamma\backslash\Lambda^+$$

where Λ^+ is the set of positive norm vectors in Λ.

One may scale the Hermitian inner product $x \bullet y$ on $\mathbb{C}^{1,9}$ so that the number of triangles is the norm of the associated vector. It turns out that Λ is an even unimodular lattice with respect to the corresponding symmetric inner product $x \cdot y := \operatorname{Re}(x \bullet y)$. However, the Hermitian form $\bullet$ is valued in $\frac{2}{3}\mathbb{Z}[\zeta_3]$, so from this perspective, it is more natural to define the rescaled Hermitian inner product $x \star y = \frac{3}{2} x \bullet y$. This choice of scaling endows Λ with the structure of an Eisenstein lattice:

Definition 2.1. An *Eisenstein lattice* is a free $\mathbb{E}$-module with an $\mathbb{E}$-valued Hermitian metric where $\mathbb{E} = \mathbb{Z}[\zeta_3]$ is the ring of the Eisenstein integers.

To avoid confusion, we use $(\Lambda, \star)$ to refer to the Eisenstein lattice and $(\Lambda^{\mathbb{Z}}, \cdot)$ to refer to the underlying integer lattice. We only use $\bullet$ to facilitate computations in the proof of Theorem 3.4. Note that $\Lambda^{\mathbb{Z}}$ is an even lattice of signature $(2, 18)$. Allcock [2], see also Proposition A.2, identifies the underlying $\mathbb{Z}$-lattice as

$$\Lambda^{\mathbb{Z}} \cong H^{\oplus 2} \oplus (-E_8)^{\oplus 2}$$

and Laza [10] verifies that $\Gamma = U(\Lambda)$ is the Hermitian isometry group of Λ. To recover the Eisenstein structure from $\Lambda^{\mathbb{Z}}$, one simply needs a fixed-point

free isometry of $\Lambda^{\mathbb{Z}}$ of order 3, corresponding to multiplication by a third root of unity.

Let S be a flat cone sphere in which μ_i singularities of deficits $2\pi/k$ have collided. The discussion after Theorem 4.1 of [17] shows the local orbifold group at $S \in \overline{\mathcal{M}}_{\alpha_1,\dots\alpha_n}$ contains a normal subgroup Γ_i of order

$$|\Gamma_i| = \frac{\mu_i!}{(1-\mu_i/k)^{\mu_i-1}}.$$

Let $\mathrm{Aut}^+(\mathcal{T})$ denote the group of oriented isomorphisms of the triangulation $\mathcal{T}$. If $\mathcal{T}$ corresponds to $v \in \Lambda^+$, we in fact have an exact sequence

$$0 \to \prod_i \Gamma_i \to \mathrm{Stab}(v) \to \mathrm{Aut}^+(\mathcal{T}) \to 0$$

where $\mathrm{Stab}(v) \subset U(\Lambda)$ is the stabilizer of the vector v. Hence we introduce a weight to each triangulation, so as to count them in the orbifold sense:

Definition 2.2. The *weight* of an oriented non-negative curvature triangulation $\mathcal{T}$ whose deficits form the partition $\mu = (\mu_1, \dots, \mu_n)$ of 12 is

$$wt(\mathcal{T}) = \frac{1}{|\mathrm{Aut}^+(\mathcal{T})|} \prod_{i=1}^{n} \frac{(1-\mu_i/6)^{\mu_i-1}}{\mu_i!} = \frac{1}{|\mathrm{Stab}(v)|}.$$

Note that the generic weight—when none of the 12 singularities have collided and $\mathcal{T}$ has no nontrivial automorphisms—is one.

Theorem 2.3. *The weighted sum of oriented non-negative curvature triangulations of S^2 with $2n$ triangles is given by the formula*

$$\sum_{|\mathcal{T}|=2n} wt(\mathcal{T}) = \frac{809}{1306069401600}\sigma_9(n).$$

Remark 2.4. Using the group of all automorphisms $\mathrm{Aut}(\mathcal{T})$ in the definition of the weight yields $\frac{809}{2612138803200}\sigma_9(n)$ for the weighted sum of unoriented triangulations, which is half the number in Theorem 2.3.

Proof of Theorem 2.3. From the above discussion and definition of the weight, we know that $\sum_{|\mathcal{T}|=2n} wt(\mathcal{T})$ is the q^n coefficient of

$$\sum_{\substack{[v]\in\Gamma\backslash\Lambda^+ \\ v\cdot v>0}} \frac{1}{|\mathrm{Stab}(v)|} q^{\frac{1}{2}v\cdot v}.$$

Since Λ is an $\mathbb{E}$-lattice whose underlying $\mathbb{Z}$-lattice is even unimodular, we can apply Corollary 3.5 of Section 3 to conclude that after adding a constant, the above series is a weight 10 modular form for $SL_2(\mathbb{Z})$ under the substitution $q = e^{2\pi i\tau}$. Furthermore, the space of weight 10 modular forms for $SL_2(\mathbb{Z})$ is one-dimensional over $\mathbb{C}$ and spanned by the weight 10 Eisenstein series

$$E_{10}(\tau) = 1 - 264\sum_{n>0}\sigma_9(n)q^n.$$



FIGURE 1. The non-negative curvature triangulations of S^2 with two triangles.



FIGURE 2. The non-negative curvature triangulations of S^2 with four triangles.

We conclude that there exists a constant $A \in \mathbb{Q}$ such that

$$\sum_{|\mathcal{T}|=2n} wt(\mathcal{T}) = A\sigma_9(n).$$

It remains to determine the constant A. The values of $\frac{\mu_i!}{(1-\mu_i/6)^{\mu_i-1}}$ for each of the values $\mu_i = 1, 2, 3, 4, 5$ are 1, 3, 24, 648, 155520 respectively. There are two triangulations consisting of two triangles, depicted in Figure 1. The corresponding partitions of 12 are $(2, 5, 5)$ and $(4, 4, 4)$ and the oriented automorphism groups have orders 2 and 6, respectively. Thus

$$A = \frac{1}{2 \cdot (3)(155520)^2} + \frac{1}{6 \cdot (648)^3} = \frac{809}{1306069401600} = \frac{809}{2^{15} \cdot 3^{13} \cdot 5^2}.$$

□

Example 2.5. We verify the formula in Theorem 2.3 for $n = 2$. Figure 2 depicts the non-negative curvature triangulations with four triangles. The partitions are $(3, 4, 5)$, $(3, 3, 3, 3)$, $(1, 1, 5, 5)$, and $(2, 2, 4, 4)$ and the oriented automorphism groups have orders 1, 12, 2, and 4 respectively. Then the

formula correctly states that

$$\frac{1}{1\cdot(24)(648)(155520)}+\frac{1}{12\cdot(24)^4}+\frac{1}{2\cdot(155520)^2}+\frac{1}{4\cdot(3)^2(648)^2}$$
$$=\frac{809}{1306069401600}(1+2^9).$$

3. Theta Series of Eisenstein Lattices

Let $I\!I_{m,n}$ denote the unique indefinite even unimodular lattice of signature (m,n). A point $p\in Gr^+(m,I\!I_{m,n}\otimes\mathbb{R})$ corresponds to a positive definite subspace $V^+\subset I\!I_{m,n}\otimes\mathbb{R}$ of dimension m. Let V^- denote the negative definite perpendicular space. Given any $v\in I\!I_{m,n}$ let $v^+\in V^+$ and $v^-\in V^-$ be its projections. As in Kontsevich [8], our starting point is the Siegel theta function on $Gr^+(m,I\!I_{m,n}\otimes\mathbb{R})\times\mathbb{H}$:

$$\Theta(p,\tau)=\sum_{v\in I\!I_{m,n}} q^{\frac{1}{2}v^+\cdot v^+}\overline{q}^{-\frac{1}{2}v^-\cdot v^-}=\sum_{v\in I\!I_{m,n}} q^{\frac{1}{2}v\cdot v}|q|^{-v^-\cdot v^-}$$

where $q=e^{2\pi i\tau}$. This function obeys the transformation property

$$\Theta(p,\gamma\cdot\tau)=\pm(c\tau+d)^{m/2}(c\overline{\tau}+d)^{n/2}\Theta(p,\tau)$$

for any $\gamma\in SL_2(\mathbb{Z})$. Furthermore, $\Theta(p,\tau)$ is absolutely invariant with respect to the action of $O(I\!I_{m,n})$ on $Gr^+(m,I\!I_{m,n}\otimes\mathbb{R})$, that is, the action on the variable p. It defines a *theta correspondence*—one can integrate against a function $f(\tau)$ on $\mathbb{H}$ to produce a function on $O(I\!I_{m,n})\backslash Gr^+(m,I\!I_{m,n}\otimes\mathbb{R})$ or vice versa. Much research has focused on former process, called the "Borcherds lift," which has led to beautiful product formulas for coefficients of modular forms [4].

Suppose that we enhance the $\mathbb{Z}$-lattice structure on $I\!I_{m,n}$ to a Hermitian structure over an imaginary quadratic field. For instance, suppose that Λ is an Eisenstein lattice of signature (m,n) whose underlying $\mathbb{Z}$-lattice is $I\!I_{2m,2n}$. Then $\Theta(p,\tau)$ can be restricted to the subvariety of $Gr^+(2m,I\!I_{2m,2n})$ parameterizing $2m$-dimensional subspaces of $I\!I_{2m,2n}\otimes\mathbb{R}$ which are defined over $\mathbb{C}$, i.e. to $Gr^+(m,\Lambda\otimes_{\mathbb{E}}\mathbb{C})$. In particular, when $m=1$, the theta correspondence $\Theta(p,\tau)$ defines a function on $\mathbb{CH}^n\times\mathbb{H}$ which is $U(\Lambda)$-invariant in the p variable and satisfies the transformation rule

$$\Theta(p,\gamma\cdot\tau)=(c\tau+d)(c\overline{\tau}+d)^n\Theta(p,\tau).$$

Let $\theta=\sqrt{3}i\in\mathbb{E}$. The following result of Basak [3] is an analogue of the uniqueness of even unimodular lattices of indefinite signature over $\mathbb{Z}$:

Proposition 3.1. *There is at most one Eisenstein lattice Λ of indefinite signature such that $\Lambda=\theta\Lambda^*$ where Λ^* is the dual lattice.*

Note that if Λ is such a lattice with Hermitian signature (m,n) then $\Lambda^{\mathbb{Z}}$ with intersection form as defined in Section 2 is an even unimodular lattice of signature $(2m,2n)$. Conversely, given an even unimodular lattice with

a fixed point free isometry of order 3, the resulting $\mathbb{E}$-lattice satisfies the condition of Proposition 3.1. Hence we introduce the following notion:

Definition 3.2. An Eisenstein lattice is $\mathbb{E}$-*unimodular* if $\Lambda = \theta\Lambda^*$. Equivalently, the norm of every vector in Λ is divisible by 3 and the underlying $\mathbb{Z}$-lattice $\Lambda^{\mathbb{Z}}$ is even unimodular.

We now define Maass forms:

Definition 3.3. A *Maass form* of weight $1-k$ is a real-analytic function $f(\tau)$ on $\mathbb{H}$ satisfying $f(\gamma\cdot\tau) = (c\tau+d)^{1-k}f(\tau)$ for all $\gamma \in SL_2(\mathbb{Z})$ which has a Fourier expansion of the form

$$c_0(\operatorname{Im}\tau)^k + \sum_{n<0} c^-(n)\Gamma(k, 4\pi|n|\operatorname{Im}\tau)q^n + \sum_{n\geq 0} c^+(n)q^n$$

and polynomial growth as $\operatorname{Im}\tau \to \infty$. Here $\Gamma(s,z) := \displaystyle\int_z^\infty x^{s-1}e^{-x}\,dx$ is the incomplete gamma function.

Siegel's foundational paper [16] on theta functions of indefinite Hermitian forms proves a generalization of our next theorem, cf. Satz 1. We present a self-contained proof which explicitly computes the Fourier coefficients relevant to the enumeration of triangulations:

Theorem 3.4. *Let Λ be an $\mathbb{E}$-unimodular Eisenstein lattice of signature $(1,k)$ with $k>1$. Let $\Gamma \subset U(\Lambda)$ be a finite index subgroup. Define*

$$g_{1-k}(\tau) := \frac{(Im\,\tau)^k}{6}\int_{\Gamma\backslash\mathbb{CH}^k} \Theta(p,\tau)\,dp$$

where dp is the complex hyperbolic volume form. Then $g_{1-k}(\tau)$ is a Maass form of weight $1-k$ in the sense of Definition 3.3, whose Fourier coefficients are tabulated in Table 1.

Proof. The proof follows from explicit computation of complex hyperbolic integrals. First, we may as well assume $\Gamma = U(\Lambda)$ as the integrand $\Theta(p,\tau)\,dp$ is $U(\Lambda)$-invariant, and so taking a finite index subgroup only changes $g_{1-k}(\tau)$ by a constant. We collect the terms of $\Theta(p,\tau)$ into Γ-orbits:

$$\Theta(p,\tau) = \sum_{[v]\in\Gamma\backslash\Lambda}\left(\sum_{w\in[v]}|q|^{-w^-\cdot w^-}\right)q^{\frac{1}{2}v\cdot v}.$$

Since $q^{\frac{1}{2}v\cdot v}$ is independent of p, we have

$$g_{1-k}(\tau) = \frac{(\operatorname{Im}\tau)^k}{6}\sum_{[v]\in\Gamma\backslash\Lambda}\left(\int_{\Gamma\backslash\mathbb{CH}^k}\sum_{w\in[v]}|q|^{-w^-\cdot w^-}\,dp\right)q^{\frac{1}{2}v\cdot v}.$$

By the results in Table 1, this Fourier series expansion of $g_{1-k}(\tau)$ converges absolutely on $\mathbb{H}$ with polynomial growth at the cusp, which ex post facto

justifies reversing the summation and integral. It is useful to introduce the following function on $\Gamma\backslash\Lambda \times \mathbb{H}$:

$$\begin{aligned} F([v],\tau) &:= \frac{1}{6}\int_{\Gamma\backslash\mathbb{CH}^k} \sum_{w\in[v]} |q|^{-w^-\cdot w^-}\, dp \\ &= \int_{\mathrm{Stab}(v)\backslash\mathbb{CH}^k} |q|^{-v^-\cdot v^-}\, dp \qquad \text{for } v \neq 0. \end{aligned}$$

The factor of $\frac{1}{6}$ disappears (except when $v = 0$) because the center of Γ, which acts on Λ by multiplication by sixth roots of unity, acts trivially on the fundamental domain $\Gamma\backslash\mathbb{CH}^k$. There are four possibilities for the behavior of this integral, depending on when $v^2 > 0$, $v^2 < 0$, v is isotropic, or $v = 0$.

Case $v^2 > 0$: Suppose $v \cdot v = 2n > 0$. Then $\mathrm{Stab}(v)$ is a finite group, and thus we can rewrite

$$\begin{aligned} F([v],\tau) &= \frac{1}{|\mathrm{Stab}(v)|}\int_{\mathbb{CH}^k} |q|^{-v^-\cdot v^-}\, dp \\ &= \frac{1}{|\mathrm{Stab}(v)|}\int_{\mathbb{CH}^k} e^{4\pi n \mathrm{Im}\,\tau(e_0^-\cdot e_0^-)}\, dp. \end{aligned}$$

where $e_0 = v/\sqrt{2n}$. Extend e_0 to an orthonormal basis of $(\mathbb{C}^{1,k}, \bullet)$. In this basis, the Klein model of $\mathbb{CH}^k$ is the unit ball in the plane $z_0 = 1$, i.e. the points $p = (1, z_1, \ldots, z_k)$ with $\sum |z_i|^2 < 1$. Let $r = \sqrt{|z_1|^2 + \cdots + |z_k|^2}$ denote the radial coordinate of this unit ball. The volume form of the metric whose curvature is pinched between $1/4$ and 1 is

$$dp = \frac{4^k r^{2k-1}\, dr\, d\mu}{(1-r^2)^{k+1}}$$

where $d\mu$ is the standard volume form on S^{2k-1}. To express the projections e_0^+ and e_0^- in terms of p, we need to be careful to use the hermitian inner product $\bullet$ since we are projecting to the *complex* span of p and its orthogonal complement; with this in mind, $e_0^+ = \frac{e_0 \bullet p}{p \bullet p} p$ and so

$$e_0^-\cdot e_0^- = e_0\cdot e_0 - e_0^+\cdot e_0^+ = 1 - \frac{|e_0 \bullet p|^2}{p\cdot p} = -\frac{r^2}{1-r^2}.$$

Let $C := 4\pi n \mathrm{Im}\,\tau$. Then, we have

$$\begin{aligned} F([v],\tau) &= \frac{4^k \mathrm{Vol}(S^{2k-1})}{|\mathrm{Stab}(v)|} \int_0^1 e^{-C\frac{r^2}{1-r^2}} \frac{r^{2k-1}}{(1-r^2)^{k+1}}\, dr \\ &= \frac{4^k \mathrm{Vol}(S^{2k-1})}{2C^k |\mathrm{Stab}(v)|} \int_0^\infty e^{-u} u^{k-1}\, du \qquad \text{where } u = C\frac{r^2}{1-r^2} \\ &= \frac{4^k (k-1)! \mathrm{Vol}(S^{2k-1})}{2C^k |\mathrm{Stab}(v)|} \\ &= \frac{(n\mathrm{Im}\,\tau)^{-k}}{|\mathrm{Stab}(v)|}. \end{aligned}$$

Case $v^2 < 0$: Suppose that $v \cdot v = -2n < 0$. We have

$$F([v],\tau) = \int_{\mathrm{Stab}(v)\backslash \mathbb{CH}^k} e^{4\pi n\, \mathrm{Im}\tau (e_1^- \cdot e_1^-)} dp$$

where $e_1 = v/\sqrt{2n}$. As before, extend e_1 to an orthonormal basis of $(\mathbb{C}^{1,k}, \bullet)$. Let $p = (1, z_1, \ldots, z_k)$ be an element of $\mathbb{CH}^k$ and let $z' = (z_2, \ldots, z_k)$. Then

$$e_1^- \cdot e_1^- = -1 - \frac{|e_1 \bullet p|^2}{p \cdot p} = -\frac{1-|z'|^2}{1-|z_1|^2-|z'|^2}.$$

We may re-write $F([v],\tau)$ as

$$\int_{\mathrm{Stab}(v)\backslash \mathbb{CH}^k} \exp\left(-4\pi n\, \mathrm{Im}\tau \frac{1-|z'|^2}{1-|z_1|^2-|z'|^2}\right) \frac{4^k\, dV_{2k}}{(1-|z_1|^2-|z'|^2)^{k+1}}$$

where dV_{2k} is the Euclidean volume form on the unit ball in $z_0 = 1$.

Since $\mathrm{Stab}(v)$ preserves the z_1 coordinate, it acts on the totally geodesic copy of $\mathbb{CH}^{k-1}$ on which $z_1 = 0$. Let X_v be a fundamental domain for this action. Then the set of $p = (1, z_1, z')$ such that $z' \in X_v$ is a fundamental domain for $\mathrm{Stab}(v)\backslash \mathbb{CH}^k$. To simplify the notation let $C := 4\pi n\, \mathrm{Im}\tau$ and $h(z') := \sqrt{1-|z'|^2}$. Then

$$\begin{aligned} F([v],\tau) &= 4^k \int_{X_v} \left(\int_0^{2\pi} \int_0^h e^{-C\frac{h^2}{h^2-r^2}} \frac{r\, dr\, d\theta}{(h^2-r^2)^{k+1}} \right) dV_{2k-2} \\ &= 4^k \int_{X_v} \left(\pi C^{-k} h^{-2k} \int_C^\infty u^{k-1} e^{-u}\, du \right) dV_{2k-2} \\ &\qquad\qquad \text{where } u = C\frac{h^2}{h^2-r^2} \\ &= 4\pi C^{-k} \Gamma(k, C) \int_{X_n} \frac{4^{k-1} dV_{2k-2}}{(1-|z'|)^k} \\ &= (n\mathrm{Im}\,\tau)^{-k} \Gamma(k, 4\pi n\, \mathrm{Im}\,\tau) \frac{\mathrm{Vol}(X_v)}{(4\pi)^{k-1}} \end{aligned}$$

Since k is a positive integer, the incomplete gamma function has the simple form $\Gamma(k, C) = (k-1)! e^{-C} \sum_{j=0}^{k-1} \frac{C^j}{j!}$.

Case $v^2 = 0$, $v \neq 0$: Let $v = \beta e$ for some $\beta \in \mathbb{Z}[\zeta_3]$ and e primitive isotropic. Since $\Lambda = \theta\Lambda^*$, an easy argument shows that there is an isotropic vector f such that $e \star f = \theta$. So $H_{\mathbb{E}} := \mathbb{E}e \oplus \mathbb{E}f$ generates a copy of the unique $\mathbb{E}$-unimodular lattice of signature $(1,1)$. Then $H_{\mathbb{E}}^{\perp}$ is a negative definite $\mathbb{E}$-unimodular lattice. Elements of $\mathrm{Stab}(v)$ correspond to triples (A, w_0, m) where A is an $\mathbb{E}$-isometry of $H_{\mathbb{E}}^{\perp}$, $w_0 \in H_{\mathbb{E}}^{\perp}$, and $m \in \mathbb{Z}$ through the action

$$\begin{aligned} e &\mapsto e \\ f &\mapsto f + \left(\frac{w_0 \star w_0}{3}\zeta_3 + m\right) e + w_0 \\ w &\mapsto -\theta^{-1}(Aw \star w_0)e + Aw \qquad \text{for } w \in H_{\mathbb{E}}^{\perp}. \end{aligned}$$

Let $\{e_2, \ldots, e_k\}$ be an orthonormal basis of $H_{\mathbb{E}}^{\perp} \otimes \mathbb{C}$ and let $\tilde{e} = -\frac{\theta}{2}e$ so that $\tilde{e} \cdot f = 1$. Let $(x, y, z_2, \ldots, z_k)$ be coordinates for $\mathbb{C}^{1,k}$ in the basis $\{\tilde{e}, f, e_2, \ldots, e_k\}$ and let $z' = (z_2, \ldots, z_k)$. Setting $y = 1$, the tuples (x, z') such that $2\mathrm{Re}(x) - |z'|^2 > 0$ give coordinates on $\mathbb{CH}^k$. Since $v^+ = \frac{v \bullet p}{p \bullet p} p$ and $v \bullet p = \frac{2}{3} v \star p = \frac{2}{3}\beta\theta$, we have

$$v^- \cdot v^- = -v^+ \cdot v^+ = \frac{-4\beta\overline{\beta}}{3(2\,\mathrm{Re}(x) - |z'|^2)}.$$

We wish to compute the following integral:

$$F([\beta e_0], \tau) = \int_{\mathrm{Stab}(v)\backslash \mathbb{CH}^k} \exp\left(-\frac{8\pi\beta\overline{\beta}\,\mathrm{Im}\,\tau}{3(2\,\mathrm{Re}(x) - |z'|^2)}\right) \frac{4^k\, dV_{2k}}{(2\,\mathrm{Re}(x) - |z'|^2)^{k+1}}.$$

Let $u = \frac{1}{2\,\mathrm{Re}(x) - |z'|^2}$ and $v = \mathrm{Im}(x)$. We have

$$\frac{-2\, dV_{2k}}{(2\,\mathrm{Re}(x) - |z'|^2)^2} = du\, dv\, dV_{2k-2}.$$

Let $\Gamma_0 \subset \mathrm{Stab}(v)$ be the finite index subgroup such that $A = Id$. We may enlarge the domain of integration to $\Gamma_0\backslash\mathbb{CH}^k$ and divide by the index $[\mathrm{Stab}(v) : \Gamma_0] = |\mathrm{Aut}(H_{\mathbb{E}}^{\perp})|$. The subgroup Γ_0 acts by translations in $H_{\mathbb{E}}^{\perp}$ on the z' coordinate and by translations by $2/\sqrt{3}$ on the v coordinate.

$$\begin{aligned} F([\beta e_0], \tau) &= \frac{4^k}{2|\mathrm{Aut}(H_{\mathbb{E}}^{\perp})|} \int_{H_{\mathbb{E}}^{\perp}\backslash \mathbb{C}^{k-1}} \int_0^{2/\sqrt{3}} \int_0^{\infty} e^{-\frac{8}{3}\pi\beta\overline{\beta}\mathrm{Im}\,\tau\, u} u^{k-1}\, du\, dv\, dV_{2k-2} \\ &= \frac{4^k (k-1)! (\frac{8}{3}\pi\beta\overline{\beta}\,\mathrm{Im}\,\tau)^{-k} \mathrm{CoVol}(H_{\mathbb{E}}^{\perp})}{\sqrt{3}|\mathrm{Aut}(H_{\mathbb{E}}^{\perp})|} \\ &= \left(\frac{2\pi\beta\overline{\beta}\,\mathrm{Im}\,\tau}{3}\right)^{-k} \frac{(k-1)!}{\sqrt{3}|\mathrm{Aut}(H_{\mathbb{E}}^{\perp})|}. \end{aligned}$$

Note that $\mathrm{CoVol}(H_{\mathbb{E}}^{\perp}) = 1$ because $((H_{\mathbb{E}}^{\perp})^{\mathbb{Z}}, \cdot)$ is unimodular.

Case $v = 0$: Immediately from the definition, $F([0], \tau) = \frac{1}{6}\mathrm{Vol}(\Gamma\backslash\mathbb{CH}^k)$. We have thus computed $F([v], \tau)$ for all $[v] \in \Gamma\backslash\Lambda$. The results are summarized in Table 1.

Case	$(\operatorname{Im}\tau)^k F([v],\tau)$
$v=0$	$\frac{1}{6}\operatorname{Vol}(\Gamma\backslash\mathbb{CH}^k)(\operatorname{Im}\tau)^k$
$v\cdot v=-2n<0$	$n^{-k}\Gamma(k,4\pi n\operatorname{Im}\tau)\dfrac{\operatorname{Vol}(X_v)}{(4\pi)^{k-1}}$
$v=\beta e$ for e primitive isotropic	$\left(\dfrac{2\pi\beta\overline{\beta}}{3}\right)^{-k}\dfrac{(k-1)!}{\sqrt{3}\lvert\operatorname{Aut}(H_{\mathbb{E}}^{\perp})\rvert}$
$v\cdot v=2n>0$	$\dfrac{n^{-k}}{\lvert\operatorname{Stab}(v)\rvert}$

TABLE 1. The coefficients of $g_{1-k}(\tau)$

We now describe the Γ-orbits of primitive isotropic vectors e. As above, choose an isotropic vector f such that $e\star f=\theta$. The $\mathbb{E}$-unimodular lattice $\{e,f\}^{\perp}$ projects isomorphically to $e^{\perp}/e$, and hence its isomorphism type is independent of f. If e_1 and e_2 lie in the same Γ orbit, then $e_1^{\perp}/e_1\cong e_2^{\perp}/e_2$. Conversely, if $e_1^{\perp}/e_1\cong e_2^{\perp}/e_2$, we may choose primitive isotropic f_1 and f_2 such that $e_i\star f_i=\theta$ and define an isometry of Λ which sends $e_1\mapsto e_2$, $f_1\mapsto f_2$, and $\{e_1,f_1\}^{\perp}\to\{e_2,f_2\}^{\perp}$. On the other hand, by Proposition 3.1, every negative definite $\mathbb{E}$-unimodular lattice L of rank $k-1$ embeds into Λ because $\Lambda\cong L\oplus H_{\mathbb{E}}$. Therefore orbits of primitive isotropic vectors in Λ, which correspond to cusps of $\Gamma\backslash\mathbb{CH}^k$, are in bijective correspondence with negative definite $\mathbb{E}$-unimodular lattices of rank $k-1$. Let

$$m_{k-1}:=\sum_{L}\frac{1}{\lvert\operatorname{Aut}(L)\rvert}$$

be the mass of negative definite $\mathbb{E}$-unimodular lattices of rank $k-1$, which is expressed in [7] in terms of generalized Bernoulli numbers.

Combining the results in Table 1 with the above observations about orbits of isotropic vectors gives

$$\begin{aligned}g_{1-k}(\tau)=&\frac{1}{6}\operatorname{Vol}(\Gamma\backslash\mathbb{CH}^k)(\operatorname{Im}\tau)^k\\&+\sum_{n<0}|n|^{-k}\,\Gamma(k,4\pi|n|\operatorname{Im}\tau)\sum_{\substack{[v]\in\Gamma\backslash\Lambda\\ v^2=2n}}\frac{\operatorname{Vol}(X_v)}{(4\pi)^{k-1}}q^n\\&+\left(\frac{3}{2\pi}\right)^k\frac{(k-1)!\zeta_{\mathbb{E}}(k)}{\sqrt{3}}\;m_{k-1}\\&+\sum_{n>0}\sum_{\substack{[v]\in\Gamma\backslash\Lambda\\ v^2=2n}}\frac{n^{-k}}{\lvert\operatorname{Stab}(v)\rvert}q^n\end{aligned}$$

where $\zeta_{\mathbb{E}}(k) = \sum_{(\beta)\neq(0)}(\beta\overline{\beta})^{-k}$ is the Dedekind zeta function. The four terms correspond to $v = 0$, $v^2 < 0$, v isotropic, and $v^2 > 0$. Since

$$\mathrm{Im}(\gamma\cdot\tau) = (c\tau+d)^{-1}(c\overline{\tau}+d)^{-1}\mathrm{Im}\,\tau,$$

we see from the transformation rule for $\Theta(p,\tau)$ that

$$g_{1-k}(\gamma\cdot\tau) = (c\tau+d)^{1-k}g_{1-k}(\tau).$$

Since $g_{1-k}(\tau)$ has the desired Fourier expansion, we conclude that $g_{1-k}(\tau)$ is a harmonic Maass form of weight $1-k$ for $SL_2(\mathbb{Z})$. □

Corollary 3.5. *Let Λ be an $\mathbb{E}$-unimodular lattice of signature $(1,k)$ and let $\Gamma\subset U(\Lambda)$ be a finite index subgroup. Then*

$$\left(\frac{1}{2\pi i}\partial_\tau\right)^k g_{1-k}(\tau) = -\frac{k!\mathrm{Vol}(\Gamma\backslash\mathbb{CH}^k)}{6(4\pi)^k} + \sum_{\substack{[v]\in\Gamma\backslash\Lambda\\ v\cdot v>0}}\frac{1}{|\mathrm{Stab}(v)|}q^{\frac{1}{2}v\cdot v}$$

is a weight $1+k$ modular form for $SL_2(\mathbb{Z})$—in fact, a scalar multiple of the Eisenstein series $E_{1+k}(\tau)$, see Proposition 3.7.

Proof. Let $D = \frac{1}{2\pi i}\partial_\tau = q\partial_q$. If $f(\tau)$ is a Maass form of weight $1-k$ with $k>1$ then by [5]

$$D^k f = (-1)^k k!\frac{c_0}{(4\pi)^k} + \sum_{n>0}c^+(n)n^k q^n$$

is a holomorphic modular form of weight $1+k$. In our case, $(-1)^k = -1$ because $k\equiv 1 \pmod 4$. The proposition follows. □

Remark 3.6. When $k=9$, Corollary 3.5 and Theorem 2.3 allow one to determine that

$$\mathrm{Vol}(\Gamma\backslash\mathbb{CH}^9) = \frac{6(4\pi)^9}{9!}\cdot\frac{1}{264}\cdot\frac{809}{2^{15}\cdot 3^{13}\cdot 5^2}.$$

This is exactly the volume computed in [12], which serves as a check for the formula in Theorem 2.3.

Proposition 3.7. *Let Λ denote the unique $\mathbb{E}$-unimodular lattice of signature $(1,k)$ and let $\Gamma = U(\Lambda)$. Assume $k>1$. Then*

$$\mathrm{Vol}(\Gamma\backslash\mathbb{CH}^k) = \frac{6(2\pi)^k}{k!(k+1)!}\prod_{j=1}^{(k+1)/2}|b_{2j}\cdot b_{2j-1,\chi}|,$$

where b_r is the Bernoulli number and $b_{r,\chi}$ is the generalized Bernoulli number associated to the quadratic character χ. The weighted number of Γ orbits of vectors of norm $2n$ is

$$\sum_{\substack{[v]\in\Gamma\backslash\Lambda\\ v\cdot v=2n}}\frac{1}{|\mathrm{Stab}(v)|} = \frac{\sigma_k(n)}{2^{k-1}k!}\prod_{j=0}^{(k-1)/2}|b_{2j}\cdot b_{2j+1,\chi}|.$$

Proof. The Maass form $g_{1-k}(\tau)$ lies in a one-dimensional space, by Theorem 2.2(1) of Lagarias and Rhoades [9]. An explicit generator of this space is the real analytic Eisenstein series $E_{1-k}(\tau, s)$ evaluated at $s = k$. Up to a choice of normalization,

$$E_{1-k}(\tau, k) = -\frac{2^{k-2}}{\pi} \sum_{(u,v)\neq(0,0)} \frac{(\operatorname{Im}\tau)^k}{(u\tau+v)(u\overline{\tau}+v)^k}.$$

The Fourier expansion of $E_{1-k}(\tau, k)$ is

$$-\frac{\zeta(1+k)(2\operatorname{Im}\tau)^k}{2\pi} + \zeta(k) + \sum_{n<0} \frac{\sigma_{-k}(|n|)}{(k-1)!}\Gamma(k, 4\pi|n|\operatorname{Im}\tau)q^n + \sum_{n>0}\sigma_{-k}(n)q^n.$$

Let C_{1-k} be the constant such that $g_{1-k} = C_{1-k}G_{1-k}$. Equating Fourier coefficients, we have

$$\left(\frac{3}{2\pi}\right)^k \frac{(k-1)!\zeta_{\mathbb{E}}(k,\chi)}{\sqrt{3}}\, m_{k-1} = C_{1-k}\zeta(k)$$

$$\frac{\operatorname{Vol}(\Gamma\backslash\mathbb{CH}^k)}{6} = C_{1-k}\frac{2^k\zeta(1+k)}{2\pi}$$

$$\sum_{\substack{[v]\in\Gamma\backslash\Lambda \\ v\cdot v = 2n}} \frac{n^{-k}}{|\operatorname{Stab}(v)|} = C_{1-k}\sigma_{-k}(n)$$

The Dedekind zeta function admits an Artin factorization

$$\zeta_{\mathbb{E}}(k) = L(k,\chi)\zeta(k)$$

where $L(k,\chi)$ is the Dirichlet L-function associated to the quadratic character mod 3. There is an explicit formula

$$L(k,\chi) = \left(\frac{\pi}{3}\right)^k \frac{2^{k-1}\sqrt{3}\, b_{k,\chi}}{k!},$$

where $b_{k,\chi}$ is the generalized Bernoulli number associated to the quadratic character χ. We have from [7] the formula

$$m_{k-1} = \frac{1}{2^{k-2}\cdot(k-1)!} \prod_{j=1}^{(k-1)/2} |b_{2j}\cdot b_{2j-1,\chi}|.$$

Using the first equation allows us to solve for the constant C_{1-k}. Next, we have the famous formula

$$\zeta(1+k) = \frac{b_{1+k}(2\pi)^{1+k}}{2(k+1)!}$$

where b_{1+k} is the Bernoulli number. Substituting the values for C_{1-k} and $\zeta(1+k)$ into the second formula gives the expression for $\operatorname{Vol}(\Gamma\backslash\mathbb{CH}^k)$, while substituting C_{1-k} into the third formula determines the weighted count of Γ-orbits of vectors of norm $2n$. □

By equating the Fourier coefficients for q^{-n}, we also get a formula for the sum of volumes of X_v over the set of orbits of vectors v of norm $-2n$.

Appendix A. The intersection pairing

Let M be a flat cone sphere with cone angle deficits $\alpha_1, \ldots, \alpha_n$. Fix a singularity $p_i \in M$. To define charts on $\mathcal{M}_{\alpha_1,\ldots,\alpha_n}$, consider a shortest path γ_j connecting p_i to p_j, for each other singularity p_j. Thurston [17] proves that γ_j are geodesics which intersect only at the endpoint p_i. Then $M - \bigcup \gamma_j$ is homeomorphic to a disk and admits an oriented, isometric embedding into $\mathbb{R}^2$ whose image is the interior of a polygon $\mathcal{P}_M$. Each path γ_j corresponds to a pair of adjacent edges v_j, w_j of $\mathcal{P}_M$ which can be glued by an oriented isometry of $\mathbb{R}^2$ to reproduce M. Relabeling the singularities if necessary, let $v_1, w_1, \ldots, v_{n-1}, w_{n-1}$ be the sequence of edges of $\mathcal{P}_M$, ordered counterclockwise, see Figure 3. Then each edge v_j or w_j is a vector in $\mathbb{R}^2$ which can be viewed as an element of $\mathbb{C}$. The following equations are satisfied:

(1) $w_j = -e^{i\alpha_j} v_j$, and
(2) $\sum_j (v_j + w_j) = 0$.

Equation (1) holds because the cone angle deficit at p_j is α_j, hence v_j and w_j meet at a vertex of $\mathcal{P}_M$ with exterior angle α_j. Equation (2) holds because the cyclically ordered sequence of edges must close up to form $\mathcal{P}_M$. One can then deform the v_j and w_j subject to equations (1) and (2) to produce a local chart $\mathcal{M}_{\alpha_1,\ldots,\alpha_n} \to \mathbb{CP}^{n-3}$, which sends

$$M \mapsto [v_1 : \cdots : v_{n-2}] \in \mathbb{CP}^{n-3}.$$

The appropriately normalized area of M is a quadratic form in $(v_1, \ldots, v_{n-2})$ which extends naturally to a Hermitian form $\bullet$ of signature $(1, n-3)$. Since M has positive area, the image of this period map actually lies in $\mathbb{CH}^{n-3}$.

When the completed moduli space is an orbifold, the developing map of the complex hyperbolic structure is a diffeomorphism from the orbifold universal cover of $\overline{\mathcal{M}}_{\alpha_2,\ldots,\alpha_n}$ to $\mathbb{CH}^{n-3}$. If the monodromy group Γ is arithmetic, then it fixes a lattice $(\Lambda, \bullet) \subset \mathbb{C}^{1,n-3}$. In certain cases, the lattice Λ can be identified with relative ease. Consider the completed moduli space $\overline{\mathcal{M}}_{\frac{2\pi}{k},\ldots,\frac{2\pi}{k}}$ of flat cone spheres modulo scaling with $2k$ singularities of cone angle deficits $\frac{2\pi}{k}$. Suppose $k = 3$, 4 or 6. Then, the moduli space is an arithmetic orbifold

$$\Gamma \backslash \mathbb{CH}^{2k-3} = \overline{\mathcal{M}}_{\frac{2\pi}{k},\ldots,\frac{2\pi}{k}}.$$

The special feature of these cases is that $\mathbb{Z}[\zeta_k]$ is a discrete subgroup of $\mathbb{C}$, and the lattice Λ can be identified in the above coordinates as the set

$$\{(v_1, \ldots, v_{2k-1}) \in \mathbb{Z}[\zeta_k]^{2k-1} \,\big|\, v_1 + \cdots + v_{2k-1} = 0\}.$$

Remark A.1. There is a slight subtlety in the construction of Λ, as the polygon $\mathcal{P}_M$ is only canonical up to rotation, but the condition



FIGURE 3. A flat cone sphere admitting a squarulation with cone angle deficits $2\pi/k$ in which v_i and w_i are identified by a rotation. Here $k = 4$.

$(v_1, \dots, v_{2k-1}) \in \mathbb{Z}[\zeta_k]^{2k-1}$ is not rotation-invariant. There are two similar approaches to rectifying this issue. First, one may define Λ as above in a single coordinate chart on $\mathcal{M}_{\frac{2\pi}{k},\dots,\frac{2\pi}{k}}$, and check that the monodromy group of the moduli space lifts to an arithmetic subgroup $\Gamma \subset U(1, 2k-3)$ preserving Λ. Alternatively, one may rigidify the embedding of $M - \bigcup \gamma_j$ into $\mathbb{R}^2$ by considering extra data on M. Let

$$\Xi \subset T(M \backslash \{p_1, \dots, p_{2k}\})$$

denote a flat sub-lattice of the tangent bundle of the smooth locus of M which is isometric at every point to $\mathbb{Z}[\zeta_k]$ with the standard metric. Then we may consider the moduli space of pairs (M, Ξ) and require that the embedding $M - \bigcup \gamma_j \to \mathbb{C}$ send Ξ_p to a translate of $\mathbb{Z}[\zeta_k] \subset \mathbb{C}$. This kills rotations of $\mathcal{P}_M$ which are not by multiples of $2\pi/k$.

The orbits $\Gamma \backslash \Lambda^+$ of positive norm vectors then correspond to pairs (M, Ξ) such that $\mathcal{P}_M$ has vertices in a translate of $\mathbb{Z}[\zeta_k]$. Thus, when $k = 3$ or 6, the elements of Λ^+ correspond to triangulations—we take a tiling $\mathbb{C}$ by equilateral triangles and intersect with $\mathcal{P}_M$ to produce a tiling of M. Similarly, when $k = 4$, the elements of Λ^+ correspond to non-negative curvature *squarulations*—decompositions of S^2 into squares such that no vertex has more than 4 squares around it, see Figure 3.

We now explicitly compute the normalized area of M as a function of $(v_1 \dots, v_{2k-1})$, so as to determine $\bullet$. A triangle with two adjacent edges

$z_1, z_2 \in \mathbb{C}$ has area $-\frac{1}{2}\mathrm{Im}(z_1\overline{z}_2)$, but to be compatible with the notation in Section 2, we normalize by $2(\mathrm{Im}\,\zeta_k)^{-1}$. The area of $\mathcal{P}_M$ is the sum of the signed areas of triangles with the following pairs of edges:

$$\begin{array}{l}(v_1,\, w_1),\\ (v_1+w_1,\, v_2),\\ \dots\\ (v_1+w_1+\cdots+v_{2k-2}+w_{2k-2},\, v_{2k-1})\\ (v_1+w_1+\cdots+v_{2k-2}+w_{2k-2}+v_{2k-1},\, w_{2k-1}).\end{array}$$

Though the last triangle has area zero, including it simplifies the formulae. Let $\beta_k = |1-\zeta_k|^2$ and $v = (v_1, \dots, v_{2k-1})$. Substituting $w_i = -\zeta_k v_i$ and adding the areas of the above triangles shows that the Hermitian norm on Λ is $\mathrm{Im}(vMv^\dagger)$, where

$$M = \frac{1}{\mathrm{Im}\,\zeta_k}\begin{pmatrix} \zeta_k^{-1} & \beta_k & \beta_k & \dots & \beta_k \\ 0 & \zeta_k^{-1} & \beta_k & \dots & \beta_k \\ 0 & 0 & \zeta_k^{-1} & \dots & \beta_k \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & \zeta_k^{-1} \end{pmatrix}.$$

Then the Hermitian product is given by

$$v \bullet w = v\frac{M - M^\dagger}{2i}w^\dagger.$$

We must restrict to the vectors v for which $v_1 + \cdots + v_{2k-1} = 0$ to compute the Hermitian form on Λ:

Proposition A.2. *Define Λ as above. Let $\epsilon_k := 2/(1+\zeta_k)$. The Gram matrix of Λ in the basis $\{e_1 - e_2, \dots, e_{2k-2} - e_{2k-1}\}$ is*

$$\begin{pmatrix} -2 & \epsilon_k & 0 & \dots & 0 \\ \overline{\epsilon}_k & -2 & \epsilon_k & \dots & 0 \\ 0 & \overline{\epsilon}_k & -2 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & -2 \end{pmatrix}.$$

Proof. Use the matrix $(M - M^\dagger)/2i$ to compute the Hermitian inner product between $e_j - e_{j+1}$ and $e_k - e_{k+1}$ for all $1 \le j, k \le 2k-2$. □

Remark A.3. The Hermitian form for Λ bears a similarity to the Dynkin diagram for A_{2k-2}. Complex reflections across the "roots" $e_j - e_{j+1}$ generate a complex hyperbolic reflection group which is likely the full automorphism group $U(\Lambda)$. The braid group of the singularities admits a representation into the monodromy group Γ of the moduli space, and the braid switching p_j with p_{j+1} maps to this complex reflection.

Appendix B. Squarulations of S^2

A non-negative curvature *squarulation*, also called a *square-tiling*, of S^2 is a decomposition of S^2 into squares such that no vertex has more than 4 squares around it. Any non-negative curvature squarulation induces the structure of a flat cone sphere by declaring each square to be isometric to the unit square. Then, there is a stratification of moduli spaces

$$\Gamma\backslash\mathbb{CH}^5 = \coprod_{\substack{\sum \mu_i = 8 \\ \mu_i \in \{1,2,3\}}} \mathcal{M}_{\mu_1\pi/2, \ldots, \mu_n\pi/2}.$$

Similar to the case of triangulations, the squarulations of S^2 correspond to Γ-orbits of positive norm lattice points in a Hermitian lattice Λ over $\mathbb{Z}[i]$ of signature $(1,5)$, where the norm $v \bullet v$ is twice the number of squares. As before, we define a weight for each squarulation which equals the reciprocal of the order of $\mathrm{Stab}(v)$:

Definition B.1. The *weight* of a non-negative curvature squarulation $\mathcal{S}$ whose deficits form the partition $\mu = (\mu_1, \ldots, \mu_n)$ of 8 is

$$wt(\mathcal{S}) = \frac{1}{|\mathrm{Aut}^+(\mathcal{S})|} \prod_{i=1}^{n} \frac{(1-\mu_i/4)^{\mu_i - 1}}{\mu_i!}.$$

The Gram matrix of Λ is computed in Proposition A.2. Substituting $k = 4$, we have $\epsilon_k = 1 - i$. In particular, every entry is divisible by $1 + i$. Since Λ has index 2^6 in its dual, we deduce that $\Lambda = (1+i)\Lambda^*$, which is analogous to $\Lambda = \theta\Lambda^*$ over $\mathbb{Z}[\zeta_3]$. As in the unimodular case, define

$$\Theta(p, \tau) = \sum_{v \in \Lambda} q^{\frac{1}{2} v^+ \cdot v^+} \overline{q}^{-\frac{1}{2} v^- \cdot v^-}.$$

The Poisson summation formula implies that

$$\Theta\left(p, \frac{-1}{2\tau}\right) = 2^6 \tau^1 \overline{\tau}^5 \Theta(p, \tau).$$

That is, Θ is modular in τ with respect to the subgroup

$$\Gamma_0(2) = SL_2(\mathbb{Z}) \cap \left\langle \begin{pmatrix} 0 & -1 \\ 2 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right\rangle.$$

The proof of Corollary 3.5 goes through, with the modification that the factor of $\frac{1}{6}$ in front of $\mathrm{Vol}(\Gamma\backslash\mathbb{CH}^k)$ must be changed to a factor of $\frac{1}{4}$ since the unit group of $\mathbb{Z}[i]$ has order 4. We conclude that

$$-\frac{5!\mathrm{Vol}(\Gamma\backslash\mathbb{CH}^5)}{4(4\pi)^5} + \sum_{\mathcal{S}} wt(S) q^{|\mathcal{S}|}$$



FIGURE 4. The squarulation with one square.



FIGURE 5. The squarulations with two squares.

is a modular form for $\Gamma_0(2)$ of weight 6, and thus lies in a two-dimensional space spanned by $E_6(\tau)$ and $E_6(2\tau)$, where

$$E_6(\tau) = 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) q^n.$$

There is one squarulation consisting one square, whose partition is $(2, 3, 3)$ and whose oriented automorphism group has order 2, see Figure 4. There are three squarulations with two squares, with partitions $(2, 3, 3)$, $(1, 1, 3, 3)$, and $(2, 2, 2, 2)$ and automorphism groups of orders 2, 2, and 8, respectively, see Figure 5. We conclude that

$$-\frac{5!\mathrm{Vol}(\Gamma\backslash\mathbb{CH}^5)}{4(4\pi)^5} + \sum_{\mathcal{S}} wt(S) q^{|\mathcal{S}|} = -\frac{1}{73728} \cdot \frac{1}{504}(E_6(\tau) + 8E_6(2\tau)).$$

Thus, we have the following formula:

Proposition B.2. *The weighted sum of non-negative curvature squarulations of S^2 with n squares is*

$$\sum_{|\mathcal{S}|=n} wt(\mathcal{S}) = \frac{1}{73728}(\sigma_5(n) + 8\sigma_5(n/2))$$

where $\sigma_5(n/2) = 0$ when n is odd.

As in Remark 3.6, we can determine the volume of the moduli space is

$$\mathrm{Vol}(\Gamma\backslash\mathbb{CH}^5) = \frac{4(4\pi)^5}{5!} \cdot \frac{1}{73728} \cdot \frac{1}{504} \cdot 9,$$

which agrees with the result in [12].

References

[1] A. D. Alexandrov. *Convex Polyhedra.* Springer Science & Business Media, 2005.

[2] D. Allcock. *The Leech lattice and complex hyperbolic reflections.* Inventiones mathematicae, 140(2): 283-301, 2000.

[3] T. Basak. *The complex Lorentzian Leech lattice and the bimonster.* Journal of Algebra, 309(1): 32-56, 2007.

[4] R. E. Borcherds. *Automorphic forms on $O_{s+2,2}(\mathbb{R})$ and infinite products.* Inventiones mathematicae 120.1: 161-213, 1995.

[5] J. Bruinier, K. Ono, and R. C. Rhoades. *Differential operators for harmonic weak Maass forms and the vanishing of Hecke eigenvalues.* Mathematische Annalen, 342(3): 673-693, 2008.

[6] P. Deligne, and G. D. Mostow. *Monodromy of hypergeometric functions and non-lattice integral monodromy.* Publications Mathmatiques de l'IHÉS 63.1: 5-89, 1986.

[7] M. Hentschel, A. Krieg, and G. Nebe. *On the classification of lattices over $\mathbb{Q}(\sqrt{-3})$, which are even unimodular $\mathbb{Z}$-lattices.* Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg, 80(2): 183-192, Springer Berlin/Heidelberg, 2010.

[8] M. Kontsevich. *Product formulas for modular forms on $O(2,n)$.* Séminaire Bourbaki 39: 41-56, 1996.

[9] J. C. Lagarias, and R. C. Rhoades. *Polyharmonic Maass forms for $PSL(2,\mathbb{Z})$.* The Ramanujan Journal 41.1-3: 191-232, 2016.

[10] R. Laza. *Triangulations of the sphere and degenerations of K3 surfaces.* arXiv: 0809.0937, 2008.

[11] E. N. Looijenga. *Uniformization by Lauricella functions—an overview of the theory of Deligne-Mostow.* Arithmetic and geometry around hypergeometric functions: 207-244, Birkhäuser Basel, 2007.

[12] C. T. McMullen. *The Gauss-Bonnet theorem for cone manifolds and volumes of moduli spaces,* 2013.

[13] G. D. Mostow. *Generalized Picard lattices arising from half-integral conditions.* Publications Mathématiques de l'IHÉS 63: 91-106, 1986.

[14] É. Picard. *Sur une extension aux fonctions de deux variables du probleme de Riemann relatif aux fonctions hypergéométriques.* Annales scientifiques de l'École Normale Suprieure 10: 305-322, 1881.

[15] R. Schwartz. *Notes on Shapes of Polyhedra.* arXiv: 1506.07252, 2015.

[16] C. L. Siegel. *Indefinite quadratische Formen und Funktionentheorie II.* Mathematische Annalen 124.1: 364-387, 1951.

[17] W. P. Thurston. *Shapes of polyhedra and triangulations of the sphere*, Geometry and Topology monographs 1: 511-549, 1998.

(Philip Engel) Harvard University
E-mail address: engel@math.harvard.edu

(Peter Smillie) Harvard University
E-mail address: smillie@math.harvard.edu